

# Pigweeds in Tennessee

*Larry Steckel, Assistant Professor, Plant Sciences*

Amaranthus is the genus for the pigweed family. The word is derived from the Greek word amaranthus, which means "everlasting." Most growers would agree that this is a very accurate description for the pigweeds. In Tennessee, over the last decade, pigweeds have become a major weed pest in row crops and pastures



*Figure 1. Palmer amaranth.*

## History

In Tennessee, pigweeds were prevalent in large numbers and were very competitive prior to the widespread use of herbicides in cropping systems. When broadcast applications of soil-applied herbicides were widely used and followed by one or more cultivations in row crops, pigweeds of any type only occasionally escaped control and developed into a real problem for the grower. However, with the reduction in usage of soil-applied and incorporated herbicides designated to control grasses and small-seeded weeds, pigweeds escaped control more often.

Sole reliance on postemergence treatments was often not as effective as preemergence or preplant incorporated herbicides for pigweed control. One reason for this is that pigweeds can emerge from March to September in Tennessee. Postemergence treatments that lack residual will only control emerged pigweeds. Cultivation, which had always been a good backup procedure for controlling pigweeds, has all but ceased. The move toward conservation tillage created a more favorable environment for germination and growth of pigweed species in cropping systems. Moreover, planting dates for row crops, particularly corn, have been pushed up much earlier in the spring. Earlier planting dates have helped increase yields but have also extended the amount of time weed control management must be active. This is primarily due to slower shading by crops under typically cooler growing conditions.

Certain pigweed biotypes have evolved resistance to the triazine and urea herbicides like atrazine, Princep, Cotoran, Direx, etc. Pigweeds that are resistant to the triazine and urea herbicides, though present in Tennessee, are not widespread. About a decade ago, several biotypes of common waterhemp and Palmer amaranth developed resistance to the ALS enzyme-inhibiting herbicides such as Scepter, Staple, Steadfast, Classic, etc. At the time, these herbicides were used extensively in soybeans and to a lesser extent in cotton and corn. Today, most populations of waterhemp and Palmer amaranth are considered resistant to the ALS class of chemistry. More recently, some biotypes of common waterhemp in the Midwest have developed resistance to the diphenyl ether chemistry, while a population of Palmer amaranth in the Carolinas has been confirmed with resistance to the dinitroaniline herbicides like Treflan and Prowl.

Of the 865 members of this family, eight pigweeds are commonly found as weeds in Tennessee. These eight weedy pigweeds can be separated into three distinct groups broken up by their growth and flowering habit.

### **Group I: Tall, Upright Growing Pigweeds with Male and Female Flowers on the Same Plant**

The first group includes **redroot pigweed** (*Amaranthus retroflexus*), **smooth pigweed** (*Amaranthus hybridus*) and **slender amaranth** (*Amaranthus gracilis*). These have been the most common pigweeds in Tennessee row crop agriculture for the last 50 years. Redroot pigweed can be found throughout much of Tennessee, but usually is not the dominant pigweed found in cropping systems. Smooth pigweed has been the pigweed in crop production. Slender amaranth can also be a problem in row crop agriculture, though not to the extent of smooth pigweed. These three species of pigweeds have similar upright growth habits, are difficult to distinguish in seedling stage, compete very well with most crops, and have both male and female flowers on the same plant (monocious).

### **Group II: Tall, Upright Growing Pigweeds with Male and Female Flowers on Separate Plants**

The second group of pigweeds includes **common waterhemp** (*Amaranthus rudis*) and **Palmer amaranth** (*Amaranthus palmeri*). They originated in the Midwest and Southwest and are relative newcomers to Tennessee. **Tall waterhemp** (*Amaranthus tuberculatus*) is very closely related to common waterhemp and is also a member of this group. Tall waterhemp is considered to be native to the Mississippi River bottom. These three pigweed species have separate male and female plants (dioecious), have an upright but somewhat branched growth habit and are very competitive with crops. They are a very diverse group and differ widely in branching and coloration of leaves/stems within species. Common and tall waterhemp are very difficult to distinguish from each other and can be found in small numbers in West Tennessee. New genetic research shows that common and tall waterhemp are very similar. Therefore, many researchers now have simply lumped them into one species, common waterhemp. Palmer amaranth is closely related to waterhemp and has long been the dominant pigweed in the Southwest. It is the most competitive and rapidly growing species of the

weedy pigweeds. Over the last few years, Palmer amaranth has become one of the most troublesome weed problems in row crops throughout West Tennessee.

### **Group III: Prostrate Growing Pigweeds with Male and Female Flowers on the Same Plant**

The third pigweed group includes **spiny amaranth** (*Amaranthus spinosus*), **tumble pigweed** (*Amaranthus albus*) and **prostrate pigweed** (*Amaranthus blitoides*). These pigweeds are similar in that they are somewhat prostrate, usually have substantial branching and are not as competitive to row crops as the other weedy pigweeds. Of these, spiny amaranth is the most predominant weedy pigweed across Tennessee. It is the only pigweed commonly a problem in both row crops and pastures. Prostrate and tumble pigweed can be found in much of Tennessee, but are typically confined to field edges. All of these weeds can occasionally cause problems in cropping systems.

### **Management**

Pigweed control is possible with a number of available effective herbicides. The pigweed species in general can best be managed when a pre-applied herbicide is followed by a post or when a post-applied herbicide is followed by another post-applied that contains some type of residual product to control pigweeds.

Herbicides such as Dual II Magnum, Cinch, Degree, Harness, Outlook, Fultime and Frontier are in many premixes and will provide residual control of pigweed in corn. Atrazine will provide both contact and residual activity on small pigweed. The plant growth regulator herbicides like Distinct, Clarity and 2,4-D are other options in corn. Callisto is a new post-applied herbicide for corn that has very good activity on pigweed. Finally, a relatively new technology, Roundup Ready Corn, can be used. Roundup (glyphosate) is a very effective pigweed herbicide.

In soybeans Dual, Prowl and Spartan will all provide residual control of pigweed. Glyphosate is a very good control option for pigweed postemergence. The diphenyl ethers like Reflex, Blazer and Cobra will also control pigweeds post if they are 4 inches in height or smaller. Treflan is still a very good option in conventional-till soybeans.

As with soybeans, glyphosate is the backbone for weed control in cotton. Glyphosate will control pigweeds provided there is thorough spray coverage. However, it does not have residual activity and Cotoran, Caparol, Dual II, Direx and Valor, or premixes that contain these products, should be considered to acquire consistent control.

## References:

Gleason, H. A. and A. Cronquist. 1963. **Manual of vascular plants.** 280-282.

Sauer, J. 1956. **Recent migration and evolution of the dioecious amaranths.** *Evolution* 11:11-31.

Sauer, J. 1950. **The Grain Amaranths: A Survey of their History and Classification.** *Annals of the Missouri Botanical Garden* 37:561-633.

Steckel, L. E., C. L. Sprague, E. W. Stoller, G. Bollero, and L. M. Wax. 2004. **Temperature effect on germination of ten *Amaranthus* species.** *Weed Sci.* 52:217-221.

Steckel, L. E., C. L. Sprague, A. G. Hager, F. W. Simmons, and G. Bollero. 2003. **Effects of shading on common waterhemp (*Amaranthus rudis*) growth and development.** *Weed Sci.* 52:898-903.

Wax, L. M. 1995. **Pigweeds of the Midwest: distribution, importance and management.** Pages 239-242 in *Proceedings of Integrated Crop Management Conference*, Vol. 7. Ames, IA: Iowa State University.

[Weed Science.com](http://www.weedscience.com). 2003. Retrieved 2004. International Survey of Herbicide Resistant Weeds. <http://www.weedscience.org/in.asp>

Visit the UT Extension Web site at:  
<http://www.utextension.utk.edu/>

05-0155 W068

Copyright 2004 The University of Tennessee. All rights reserved. This document may be reproduced and distributed for nonprofit educational purposes providing that credit is given to University of Tennessee Extension.

Programs in agriculture and natural resources, 4-H youth development, family and consumer sciences, and resource development.  
University of Tennessee Institute of Agriculture, U.S. Department of Agriculture and county governments cooperating.  
UT Extension provides equal opportunities in programs and employment.