


Preventing Off-target Herbicide Problems in Vineyards



Trevor D. Israel, UT Extension Assistant

G. Neil Rhodes Jr., Professor and UT Extension Weed Management Specialist

Introduction

Grape production is an increasingly important component of agriculture in Tennessee and many parts of the United States. Since many vineyards are located near pastures, hay fields or roads, off-target movement of pasture and right-of-way herbicides is becoming a major issue in grape production. While these herbicides are valuable tools for weed management, off-target damage to grape often results in expensive fines and/or lawsuits, reduced yields, delayed ripening, and poor fruit quality. Fortunately, preventive steps can be taken to avoid these problems.

Herbicide Selection

Although highly effective on several broadleaf weeds in pastures and rights-of-way, the auxin, or growth regulator herbicides can damage sensitive crops if not used properly. The characteristics of these herbicides determine which product to use in different situations. Under the right conditions, volatile herbicides change from a liquid to gas or vapor and move away from the target. Typically, dicamba and 2,4-D are more volatile than aminopyralid or picloram. Keep in mind that while ester formulations are more likely to volatilize than amine (salt) formulations, some salts of 2,4-D and dicamba are more volatile than others.



*Fig. 1. Off-target herbicide damage to grape.
(Photo by Jay W. Pscheidt)*

The persistence of herbicides is also important to consider when choosing a product. While dicamba and 2,4-D are highly active on grape even in minute doses, these materials are relatively nonpersistent in soil and in treated pasture grasses and hay. However, aminopyralid and picloram can stay active in soil, pasture grass and hay for a year or longer. When these forages are consumed by animals, the chemical passes through their digestive and urinary systems without change and into the manure and urine. It takes several days for aminopyralid and picloram to pass through the digestive and urinary systems of an animal. Producers should handle treated hay and manure so that placement will not affect future production. **Herbicides that contain aminopyralid or picloram are for use in permanent grass pastures and grass hay fields only. They should not be used in fields that will be converted into a vineyard or rotated to other broadleaf crops.**

Another characteristic to consider is water solubility. Picloram is more soluble than aminopyralid and therefore more likely to be moved off-site by runoff.

Newer pasture and right-of-way herbicides such as aminopyralid have strong attributes in that they control some of our worst pasture and hay field weeds such as horsenettle, tall ironweed and beggarweed. Also, volatility, unlike in the case of 2,4-D, is not an issue.

Drift Prevention

Because vineyards are often located near pastures, hay fields and other row crops, the probability of herbicide drift is high unless preventive measures are taken. Several factors can contribute to herbicide drift to sensitive areas. Two types of drift, physical and vapor, can occur. Physical drift is the movement of liquid spray droplets away from the target, and it is influenced by spray equipment and wind. Calibrating your sprayer for low pressure (30 psi or less) and high volume (20 gallons per acre or more) applications will reduce the number of fine spray droplets. Lowering the boom height also reduces drift, but make sure that the correct spray pattern overlap is achieved. New advances in spray tip design, such as air induction technology and Turbo TeeJet Induction nozzles, have allowed for adequate spray patterns while producing large droplets. In many cases, induction nozzles require higher operating pressures than flat fan nozzles, so be sure to check the nozzle specifications. Another rule of thumb is to spray on calm days and when wind direction is away from sensitive areas. Calm conditions are more likely to occur early or late in the day. Another common sense approach is to allow some buffer area that is not sprayed adjacent to sensitive crops. Drift reduction agents can also be used to reduce physical drift, but check the labels for compatibility.

Common auxin herbicides.

Common name	Chemical family	Trade names
aminopyralid	Pyridine-carboxylic acid	Milestone, ForeFront R&P, ForeFront HL, GrazonNext
picloram	Pyridine-carboxylic acid	Tordon, Surmount, Grazon P+D
2,4-D	Phenoxyacetic acid	Various names and mixtures
dicamba	Benzoic acid	Banvel, Clarity, Oracle, Rifle, Brush, Rangestar, Weedmaster



Fig. 2. Herbicide applications made on calm days with low pressures and low boom heights can reduce the likelihood of physical drift.

Vapor drift is the movement of spray vapor away from the target after the herbicide has been deposited on the target. It is mainly influenced by the vapor pressure of the herbicide formulation, air temperature and relative humidity (RH). Ester formulations have higher vapor pressures and are therefore more volatile than amine formulations. If sensitive crops are nearby, use the amine formulation of 2,4-D rather than the low volatile ester formulation. Some chemicals volatilize readily at warm (greater than 85 F) temperatures, and dry air (RH less than 40 percent) increases the likelihood of vapor drift. During late spring to summer applications, warm temperatures are likely to be encountered at or shortly after spraying. Keep in mind that vapor drift will be worse under warm conditions, and that it can occur even a few days after application. Dicamba herbicides are also temperature sensitive (see table). Drift reduction measures such as low pressure, special nozzles, drift retardants, etc. do not reduce vapor drift.

Other considerations to bear in mind are proximity to sensitive areas and timing of herbicide application. Be familiar with adjoining properties and owners, so you know the location and growth stage of their grapevines. If you manage a vineyard, keep your neighbors updated often, especially since dormant times can vary from year to year. If grapes are nearby, try to time herbicide applications before bud break or after leaf drop. This timing is often difficult to accomplish, because the optimum time for weed control may occur when grapes are actively growing. However, some weeds, such as musk thistle and buttercup, may be treated after mid-October with 2,4-D. They also can be sprayed with 2,4-D in late winter or early spring. These are good approaches for a field across the fence from your neighbor's vineyard, in that you could treat at a time of year when the vines are dormant. A little communication with your neighbors can go a long way to reduce the likelihood of negative consequences.

Sprayer Contamination

A common way for pasture herbicides to make it into vineyards is through sprayer contamination. Pasture herbicides are notoriously difficult to rinse from sprayers. Rinsing with water does not remove all herbicide residues from a sprayer and hoses. Residues are bound to porous materials such as rubber, plastic and polyethylene and can be dissolved later when another application is made. Common solvents used for cleaning sprayers include ammonia and various commercial cleaning agents. The longer a herbicide sits in the sprayer, the more difficult it will be to clean, so always try to end the day with a clean sprayer. Stainless steel tanks can be easily cleaned, but rubber hoses and fittings should be replaced to avoid contamination. Be sure to read the herbicide label for proper cleaning instructions. Due to the difficulty with which these herbicides are removed from a sprayer, dedicating a sprayer to be used only on pastures and hay fields is the best way to avoid herbicide tank contamination issues in grapes and other sensitive crops. Do not use pasture herbicides in sprayers that may be used to manage vegetation in the vineyard.



Fig. 3. Off-target herbicide damage to grape can delay ripening and reduce yield.

Field Selection

The location, characteristics, and history of a field influence future management strategies. A proper risk assessment should be performed before spraying a pasture with some of these herbicides. Rains can wash certain herbicides downhill to off-target areas. Applications of picloram should not be made on sites with steep slopes and bare soils. It is also important to avoid situations where herbicide runoff can contaminate streams or reservoirs that may be used to irrigate grape or other sensitive crops. Vegetated buffer strips around ponds can help to reduce herbicide surface movement. Before establishing a vineyard, one should research the history of the field to see if it has been treated with any persistent herbicides in the past three years. Also, a vineyard should not be established in an area of a field where hay that has been treated with aminopyralid or picloram has been stored.

Monitoring Results

Producers are encouraged to assess the performance of herbicides in pastures and hay fields. It is important to keep a log of all applications with dates, products, field locations and weather conditions. (This log is required by law for picloram, as it is a restricted use pesticide.) Adequate records will help producers keep herbicides contained within the target area, thereby reducing negative impacts to other farm operations and insuring the availability of these important tools for the foreseeable future.

Important considerations for spraying near vineyards:

- *Proximity of sensitive crops*
- *Herbicide label instructions*
- *Potential volatility of the herbicide you choose*
- *Separate sprayer for pastures and hay fields*
- *Current and forecast weather*
- *Sprayer calibration and adjustment to minimize drift*
- *Potential runoff into sensitive areas or irrigation waters*
- *Inform your neighbors of your plans*

References

- Anonymous. 2012. Aminopyralid Stewardship Sheet. Dow AgroSciences. Available online at <http://www.dowagro.com/range/products/milestone.htm>
- Anonymous. 2012. Milestone herbicide label. Dow AgroSciences.
- Johnson, Bill, Bill Casady, Dallas Peterson, and Dennis Kuhlman. 1997. Cleaning field sprayers to avoid crop injury. University of Missouri Extension Publication G4852.
- Steckel, L., C. Craig, and A. Thompson. 2005. Cleaning plant-growth-regulator (PGR) herbicides out of field sprayers. University of Tennessee Publication W 071.

Picture Credits

Fig. 1. Grape damage. Digital image. Accessed 10 May 2013. Available online at <http://pnwhandbooks.org/plantdisease/node/6087/print>

Fig. 3. Grapes. Digital image. Accessed 10 May 2013. Available online at <http://www.tennesseewines.com/grapes.htm>

Development of this fact sheet was funded in part by a grant from Philip Morris International, with additional support from Dow AgroSciences and DuPont Crop Protection.

ag.tennessee.edu

Disclaimer

This publication contains herbicide recommendations that are subject to change at any time. The recommendations in this publication are provided only as a guide. It is always the pesticide applicator's responsibility, by law, to read and follow all current label directions for the specific herbicide being used. The label always takes precedence over the recommendations found in this publication.

Use of trade or brand names in this publication is for clarity and information; it does not imply approval of the product to the exclusion of others that may be of similar, suitable composition, nor does it guarantee or warrant the standard of the product. The author(s), the University of Tennessee Institute of Agriculture and University of Tennessee Extension assume no liability resulting from the use of these recommendations.

W297-A 07/13 14-0012

Programs in agriculture and natural resources, 4-H youth development, family and consumer sciences, and resource development.

University of Tennessee Institute of Agriculture, U.S. Department of Agriculture and county governments cooperating.

UT Extension provides equal opportunities in programs and employment.