

Management of Small Grain Diseases Fungicide Efficacy for Control of Wheat Diseases 2018

*Heather M. Kelly, Associate Professor and Extension Specialist
Department of Entomology and Plant Pathology*

The North Central Regional Committee on Management of Small Grain Diseases (NCERA-184) has developed the following information on fungicide efficacy for control of certain foliar diseases of wheat for use by the grain production industry in the U.S. Efficacy ratings for each fungicide listed in the table were determined by field testing the materials over multiple years and locations by the members of the committee. Efficacy is based on proper application timing to achieve optimum effectiveness of the fungicide as determined by labeled instructions and overall level of disease in the field at the time of application. Differences in efficacy among fungicide products were determined by direct comparisons among products in field tests and are based on a single application of the labeled rate as listed in the table. Table includes most widely marketed products, and is not intended to be a list of all labeled products. *See next page for table footnotes.*

Efficacy of fungicides for wheat disease control based on appropriate application timing.

Fungicide(s)				Powdery mildew	Stagonospora leaf/glume blotch	Septoria leaf blotch	Tan spot	Stripe rust	Leaf rust	Stem rust	Head scab	Harvest Restriction
Class	Active ingredient	Product	Rate/A (fl. oz)									
Strobilurin	Picoxystrobin 22.5%	Aproach SC	6.0 - 12	G ¹	VG	VG ²	VG	E ³	VG	VG	NL	Feekes 10.5
	Fluoxastrobin 40.3%	Evito 480 SC	2.0 - 4.0	G	--	--	VG	--	VG	--	NL	Feekes 10.5 and 40 days
	Pyraclostrobin 23.6%	Headline SC	6.0 - 9.0	G	VG ²	VG ²	E	E ³	E	G	NL	Feekes 10.5
Triazole	Metconazole 8.6%	Caramba 0.75 SL	10.0 - 17.0	VG	VG	--	VG	E	E	E	G	30 days
	Tebuconazole 38.7%	Folicur 3.6 F ⁵	4.0	NL	NL	NL	NL	E	E	E	F	30 days
	Prothioconazole 41%	Proline 480 SC	5.0 - 5.7	--	VG	VG	VG	VG	VG	VG	G	30 days
	Prothioconazole 19% Tebuconazole 19%	Prosaro 421 SC	6.5 - 8.2	G	VG	VG	VG	E	E	E	G	30 days
	Propiconazole 41.8%	Tilt 3.6 EC ^{4,5}	4.0	VG	VG	VG	VG	VG	VG	VG	P	Feekes 10.5
Mixed modes of action ⁶	Tebuconazole 22.6% Trifloxystrobin 22.6%	Absolute Maxx SC	5.0	G	VG	VG	VG	VG	E	VG	NL	35 days
	Cyproconazole 7.17% Picoxystrobin 17.94%	Aproach Prima SC	3.4 - 6.8	VG	VG	VG	VG	E	VG	--	NR	45 days
	Prothioconazole 16.0% Trifloxystrobin 13.7%	Delaro 325 SC	8.0	G	VG	VG	VG	VG	VG	VG	NL	Feekes 10.5 35 days
	Fluoxastrobin 14.8% Flutriafol 19.3%	Fortix Preemptor SC	4.0 - 6.0	--	--	VG	VG	E	VG	--	NL	Feekes 10.5 and 40 days
	Fluapyroxad 2.8% Pyraclostrobin 18.7% Propiconazole 11.7%	Nexicor EC	7.0 - 13.0	G	VG	VG	E	E	E	VG	NL	Feekes 10.5
	Fluxapyroxad 14.3% Pyraclostrobin 28.6%	Priaxor	4.0 - 8.0	G	VG	VG	E	VG	VG	G	NL	Feekes 10.5
	Propiconazole 11.7% Azoxystrobin 13.5%	Quilt Xcel 2.2 SE ⁵	10.5 - 14.0	VG	VG	VG	VG	E	E	VG	NL	Feekes 10.5
	Prothioconazole 10.8% Trifloxystrobin 32.3%	Stratego YLD	4.0	G	VG	VG	VG	VG	VG	VG	NL	Feekes 10.5 35 days
	Benzovindiflupyr 10.3% Propiconazole 11.7% Azoxystrobin 13.5%	Trivapro SE	9.4 - 13.7	VG	VG	VG	VG	E	E	VG	NL	Feekes 10.5.4
	Metconazole 7.4% Pyraclostrobin 12%	TwinLine 1.75 EC	7.0 - 9.0	G	VG	VG	E	E	E	VG	NL	Feekes 10.5

Footnotes

¹*Efficacy categories: NL = Not Labeled; NR = Not Recommended; P = Poor; F = Fair; G = Good; VG = Very Good; E = Excellent; -- = Insufficient data to make statement about efficacy of this product.*

²*Product efficacy may be reduced in areas with fungal populations that are resistant to strobilurin fungicides.*

³*Efficacy may be significantly reduced if solo strobilurin products are applied after stripe rust infection has occurred.*

⁴*Application of products containing strobilurin active ingredients may result in elevated levels of the mycotoxin Deoxynivalenol (DON) in grain damaged by head scab.*

⁵*Multiple generic products containing the same active ingredients also may be labeled in some states.*

⁶*Products with mixed modes of action generally combine triazole and strobilurin active ingredients. Nexicor, Priaxor and the Trivapro include carboxamide active ingredients.*

Disclaimer

This publication contains pesticide recommendations that are subject to change at any time. The recommendations in this publication are provided only as a guide. It is always the pesticide applicator's responsibility, by law, to read and follow all current label directions for the specific pesticide being used. The label always takes precedence over the recommendations found in this publication.

Use of trade or brand names in this publication is for clarity and information; it does not imply approval of the product to the exclusion of others that may be of similar, suitable composition, nor does it guarantee or warrant the standard of the product. The author(s), the University of Tennessee Institute of Agriculture and University of Tennessee Extension assume no liability resulting from the use of these recommendations.

AG.TENNESSEE.EDU

Real. Life. Solutions.™